Key research skills

Understanding modelled economic evaluations: a reader's guide for clinicians

conomic evaluations have a long history in health care. 1,2 Full economic evaluations aim to inform decision making through comparing the costs and outcomes of two or more interventions, strategies, programs or policies, to estimate their efficiency via an incremental cost-effectiveness ratio. The premise for conducting economic evaluations is that health care resources are finite, and there is an opportunity cost when resources are allocated to one health care intervention over another.^{3,4} In Australia, economic evaluations are important considerations in policy decisions on what should be publicly funded under the Pharmaceutical Benefits Scheme (PBS)⁵ and Medicare Benefits Schedule (MBS).⁶ Furthermore, clinician– researchers are increasingly considering both clinical effectiveness and cost-effectiveness in evaluation studies and funding applications.

Full economic evaluations are classified by the type of evaluation, with the most common types being cost-effectiveness analysis (CEA), cost-utility analysis (CUA) and cost-benefit analysis (CBA).³ The method for conducting these economic evaluations can be study-based or decision–analytic model-based, or both.⁷ Modelled economic evaluations can overcome some of the limitations associated with study-based economic evaluations.^{7,8} The complexity and use of modelled evaluations has increased with improved computing power and data availability. Several published articles offer clinicians an introduction to

economic evaluations,^{1,9,10} but few to date have focused on modelled evaluations.^{11,12} In this key research skills article, we aim to improve clinician understanding of modelled evaluations. In the Supporting Information, we illustrate key modelling concepts using two recently published models in the *Medical Journal of Australia*.

Why modelled economic evaluations?

Why are model-based evaluations done? Modelled evaluations can be performed alongside empiric studies or as standalone studies. Modelled evaluations do not replace study-based evaluations — rather, they enable evidence synthesis across multiple studies into relevant decision-making contexts, extrapolation of trial-based results beyond the time horizon, and hypothesis generation where data are unavailable. Results beyond the study-based and model-based economic evaluations differ.

We use an example of a randomised controlled trial (RCT) calculating the cost-effectiveness of a new blood pressure medicine compared with placebo to illustrate why modelled evaluations might be required. Firstly, the model can be used to synthesise evidence across multiple trials for decision making. Whereas the RCT is only comparing the new medicine against the placebo, a modelled evaluation can extend this to compare the new medication against several

	Study-based economic evaluations	Modelled economic evaluations
Description	Empiric studies include economic evaluations done alongside an RCT, and quasi- experimental studies (eg, before–after studies)	Standalone or done alongside study-based economic evaluations. Use mathematical models (decision–analytic models) to synthesise relevant evidence for decision making, extrapolate costs and benefits beyond a trial-based follow-up period, and generate hypotheses where data are unavailable
Interventions compared	Usually limited by study design (eg, only compares control and intervention arms studied in an RCT)	Can consider control versus several relevant interventions, across multiple empiric trials
Parameters (eg, probabilities, costs, outcomes)	Data collected during study (eg, within-trial measurements of costs and outcomes), although secondary data may also be used to estimate costs that are not captured within the study	Data from a variety of sources, including primary data, secondary published data, expert opinion
Time horizon	Limited by study duration	Often longer, can extrapolate costs and outcomes up to a lifetime time horizon
Uncertainty	Primary research data collected; however, sensitivity analyses are needed to characterise impact of uncertainty on results	Parameters rely on various primary and/or secondary data, which introduces additional uncertainty into ICER estimates
Generalisability	Limited, as population reflects context of study (eg, patient demographics, health care setting)	Broader generalisability, can simulate patient populations, interventions, and health care settings, and inform decision making across different contexts

Winnie Chen^{1,2} D

Martin Howell²

Alan Cass¹

Gillian Gorham¹

Kirsten Howard²

1 Menzies School of Health Research, Charles Darwin University, Darwin, NT.

2 Menzies Centre for Health Policy and Economics, University of Sydney, Sydney, NSW.

> winnie.chen@ menzies.edu.au

2 Definitions

Quality-adjusted life years (QALY)

QALY = Utility weight × years lived

QALYs are a combined measure of the quality and quantity of life lived. When QALYs are modelled as an outcome, a utility weight is assigned to each health state, where 0 represents death and 1 represents perfect health. For example, if stroke has a utility weight of 0.52 and an individual lives for 10 years, their QALY will be 5.2 QALYs.

Incremental cost-effectiveness ratio (ICER)

Cost (intervention) – cost (control)

QALY (intervention) – QALY (control)

ICER is the difference (increment) in costs, divided by the differences (increment) in effects between two treatments. ICER provides decision makers with an estimate of the additional costs required to achieve each additional outcome, that is, efficiency. The decision maker can then use these results, alongside other considerations (eg, clinical, political) to assess whether the intervention represents value for money in their context. What ICER is considered cost-effective? In the United Kingdom, there is an explicit cost-effectiveness ratio of about £20 000/QALY for most circumstances, whereas Australia does not have an explicit threshold.¹⁴

commonly used blood pressure agents not included in the trial.

Secondly, models can extrapolate findings beyond the trial-based follow-up period (time horizon). If the above RCT was conducted over two years, the quality-adjusted life years (QALYs; Box 2) gained through improved blood pressure management are likely to be accrued over an individual's lifetime (eg, long term reduction in cardiovascular events) rather than simply within the two-year follow-up period of the RCT. Therefore, the model can be used to extrapolate cost-effectiveness to an appropriate time horizon.⁷

Thirdly, using cost-effectiveness results from a single trial can be problematic if the trial is not generalisable to the population or health care setting in which the decision is being made. For example, if the RCT was done in the United States, a modelled evaluation can use Australian estimates of health care use and costs to assist with decision making on whether the new medication is cost-effective in our local context.

Which models are used and why?

The type of decision–analytic model used depends on several factors, including the decision context, data availability, and interpretability of the model. ¹⁵ Below we cover several common model types including simple decision trees, Markov cohort, and Markov microsimulation models. Box 3 summarises common terms used in modelled economic evaluation.

Simple decision tree


Consider a simple example of calculating the costeffectiveness of a new blood pressure medicine compared with placebo. A simple decision tree can be constructed with the one-year probabilities of developing outcomes of cardiovascular disease (CVD) and death. In Box 4, the intervention cohort had a lower probability of developing CVD (15%) compared with the placebo cohort (25%). Costs and outcomes are calculated for both intervention and placebo groups (see example of how cost is calculated for the new blood pressure medicine in Box 4). From costs and outcomes, an incremental cost-effectiveness ratio (ICER; Box 2) can be calculated, for example, \$30000 per QALY gained. The decision tree is easily interpretable but infrequently used because of inherent difficulties in incorporating longer term cost-effectiveness.15

Markov cohort


Markov models are widely used to represent the movement between different health states over time. Box 5 shows a Markov cohort model for a similar scenario as described above. For simplicity, we assume that the annual probabilities of developing outcomes of CVD and death remain constant over 30 years. If we model a population of 100 patients in a Markov cohort model, the entire cohort enters and moves through the model for 30 years, at one-year cycles. All patients have no CVD at the start, but by the end of Year 1, 15 (15% of 100 people) with no CVD develop CVD. Box 6 follows the cohort through Years 1 to 4. The ICER now reflects the differences between intervention and control groups over 30 years, rather

3 Common terms used in modelled economic evaluations

Term	Definition and examples ates Different states of health included in the model eg, no cardiovascular disease, cardiovascular disease death.	
Health states		
Cycle length	Time per cycle in a Markov model eg, commonly one year but can vary.	
Time horizon	Follow-up period in the modelled analysis eg, 30 years (30 cycles, if cycle length is one year).	
Parameters	Input data including transition probabilities, costs and outcomes.	
Transition probability	Probability of moving from one health state to another, within one cycle in a Markov model.	
First- versus second-order Monte Carlo simulation	First-order Monte Carlo simulation refers to microsimulation, or individual "walk-through" eg, in Markov microsimulation models. Second-order Monte Carlo simulation refers to probabilistic sensitivity analysis (PSA).	
One-way, two-way, multi-way or probabilistic sensitivity analysis	One-way sensitivity analysis examines changes in cost-effectiveness when one parameter is varied. Two-way and multi-way sensitivity analyses examine changes when two or more parameters are varied. PSA examines the effect of changing multiple parameters simultaneously through sampling from distributions.	



BP = blood pressure; CVD = cardiovascular disease. A simple decision tree constructed in TreeAge Pro. Blue box represents decision node, green circle represents chance node, red triangle represents terminal state. Example calculation of cost in the "New BP medicine" group: total cost (new BP medicine group) = 0.8*800 + 0.15*5800 + 0.05*6800 = \$1850.




BP = blood pressure. A Markov model constructed in TreeAge Pro (simplified representation). Blue box represents decision node, purple circle represents Markov node, green circle represents chance node, red triangle represents terminal state.

than the one year as per the simple decision tree. To conduct the same analysis using a simple decision tree would require an additional branch for each of the 30 years modelled.

Markov microsimulation

For the same scenario as above, a Markov microsimulation model, also known as a first-order Monte Carlo simulation, can be constructed. The structure of the model would be the same as the Markov cohort model (Box 5). However, in microsimulation, each of the 100 patients moves

through the model individually until the end of 30 years. For example, in Box 6, Person 1 develops CVD in Year 4 (orange bubble), and the model follows this individual's journey over 30 years or until death. The simulation is repeated for each person until all 100 individuals complete their 30 years simulated disease journey. Microsimulation enables greater model complexity. For example, the probability of death for Person 1 can be changed to reflect their individual demographic profile (eg, age, sex), or their number of years living with CVD. Using microsimulation, costeffectiveness for subpopulations can be calculated separately — for example, separate ICER of new



CVD = cardiovascular disease. Visual representation of Markov cohort (left) versus Markov microsimulation (right) in initial Years 1 to 4. In the Markov cohort model, the smiley faces represent the group of individuals moving through the model, according to probabilities of 0 to 1 (purple for probabilities from no CVD to other outcomes; orange for probabilities from CVD; blue for probabilities from death). The numbers within the bubbles represent the number of individuals in each health state at the end of each year. In the Markov microsimulation model, the smiley face represents a single individual moving through the model. The orange bubbles represent the pathway this individual takes through the simulation from Years 1 to 4. In the microsimulation, the individual simulations are repeated for a number of patients (eg, 100 individuals).

medicine versus placebo for male and female patients.

Discrete event simulation


Discrete event simulation (DES) is a microsimulation model where individuals move through health states according to time-to-event probability distributions. These are less commonly used in economic evaluations than above models, but may offer greater flexibility than Markov microsimulation, for example, in its treatment of time and handling of the simultaneous risk of multiple events. ^{16,17}

Model uncertainty and sensitivity analysis

A model should not only provide results on the incremental cost-effectiveness of an intervention, but also address the question of which assumptions would result in a higher or lower ICER. Uncertainty of cost-effectiveness results due to uncertainty in model assumptions is unavoidable. 8,13,18 Uncertainty can be broadly categorised into methodological, structural, and parameter uncertainty. Methodological

uncertainty refers to methodological selection of model parameters, such as decisions on time horizon or cycle length. Structural uncertainty refers to uncertainty around the model structure, such as deciding which health states are chosen to reflect the disease process. Parameter uncertainty refers to uncertainty around input data, including probability of an event, costs, and health outcome estimates. For example, parameter uncertainty may present in the form of standard deviations or confidence intervals around a mean; or may arise as a result of multiple literature estimates of parameter values.

A sensitivity analysis assesses the impact of parameter uncertainty. Deterministic sensitivity analysis includes one-way, two-way, and multi-way sensitivity analyses, and these involve varying the inputs for one or more parameters, and seeing the influence those changes have on the costs, outcomes and ICERs. For example, in Box 5, the probability of death is 5% in people with CVD. The one-way sensitivity analysis can examine how ICER changes when a range of alternative probabilities from 3% to 7% are used instead of 5%. Deterministic sensitivity analysis is useful in



QALY = quality adjusted life years; WTP = willingness to pay (threshold). Generated in TreeAge Pro. The diagonal dotted line represents an arbitrary willingness to pay threshold of \$50 000 per QALY. Red dots represent each run that is above the willingness to pay threshold (considered not cost-effective), whereas green dots represent each run that is below the willingness to pay threshold (considered cost-effective). The larger green circle represents a 95% confidence interval around the estimated ICER of \$75742 per QALY.

identifying which parameters have the largest effect on costs, health outcomes, and cost-effectiveness, and whether the ICER varies from cost-effective to not costeffective with the parameter changes.

Probability sensitivity analysis (PSA) involves varying one or more parameters by entering them as a distribution, rather than as a single fixed value. 19 This is also known as second-order Monte Carlo simulation. If we conduct a PSA for the Markov cohort model in Box 5, the parameters (probabilities, costs, outcomes) are entered as distributions rather than fixed values. For example, instead of entering an annual probability of death of 5% for people with CVD, we now enter the probability of death as a distribution with a mean value of 0.05. The model is then run multiple times (eg, 1000 runs) and a new value for the transition probability is selected from this distribution each time. Visually, PSA results can be plotted on a costeffectiveness plane, with the ICER from each run represented as a single dot (Box 7). The diagonal dotted line in Box 7 represents an arbitrary willingness to pay threshold of \$50000 per QALY. Red dots represent each run that is above the willingness to pay threshold (considered not cost-effective), whereas green dots represent each run that is below the willingness to pay threshold (considered cost-effective). The larger green circle represents a 95% confidence interval around the estimated ICER of \$75742 per QALY.

A reader's guide to understanding recently published models

We apply our understanding of modelled economic evaluations to published models, using examples

from two 2023 articles published in the *Medical Journal of Australia*. The Markov cohort model by Xiao and colleagues examines the cost-effectiveness of chronic hepatitis B screening strategies against usual care. The Markov microsimulation model by Venkataraman and colleagues examines the cost-effectiveness of several risk score and coronary artery calcium score-based strategies for initiating statin therapy. We have summarised key information from these two modelled evaluations in the Supporting Information, table. This table seeks to highlight key concepts rather than apply a relevant critical appraisal tool such as CHEERS (Consolidated Health Economic Evaluation Reporting Standards) or similar. 22-24

In conclusion, understanding modelled economic evaluations is valuable for clinicians involved in health research or policy decisions. We encourage readers interested in health economics to access in-depth resources, which include worked examples on how to construct a model. ^{4,8,15}

Open access: Open access publishing facilitated by Charles Darwin University, as part of the Wiley - Charles Darwin University agreement via the Council of Australian University Librarians.

Competing interests: No relevant disclosures.

Provenance: Not commissioned; externally peer reviewed.

© 2024 The Author(s). *Medical Journal of Australia* published by John Wiley & Sons Australia, Ltd on behalf of AMPCo Pty Ltd.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

- 1 Evans DB. What is cost-effectiveness analysis? *Med J Aust* 1990; 153 Suppl 1: 7-9.
- 2 Hurley S. A review of cost-effectiveness analyses. *Med J Aust* 1990; 153 Suppl 1: 20-23.
- **3** Drummond MF, Sculpher MJ, Claxton K, et al. Methods for the economic evaluation of health care programmes. Oxford: Oxford University Press, 2015.
- 4 Gray AM. Applied methods of cost-effectiveness analysis in healthcare. United Kingdom: Oxford University Press, 2010.
- 5 Department of Health and Aged Care. Guidelines for preparing submissions to the Pharmaceutical Benefits Advisory Committee (PBAC). Canberra: Australian Government, 2016. https://pbac.pbs. gov.au/ (viewed July 2023).
- 6 Department of Health and Aged Care. Guidelines for preparing assessments for the MSAC: Australian Government, 2021. http:// www.msac.gov.au/internet/msac/publishing.nsf/Content/MSAC-Guidelines (viewed July 2023).
- 7 Sculpher MJ, Claxton K, Drummond M, McCabe C. Whither trial-based economic evaluation for health care decision making? Health Econ 2006; 15: 677-687.
- 8 Briggs A, Sculpher M, Claxton K. Decision modelling for health economic evaluation. Oxford: Oxford University Press, 2006.
- 9 Wong G, Howard K, Craig JC. Economic evaluation in clinical nephrology: Part 1. an introduction to conducting an economic evaluation in clinical practice. *Nephrology (Carlton)* 2010; 15: 434-440.
- **10** Goodacre S, McCabe C. An introduction to economic evaluation. *Emerg Med* / 2002; 19: 198-201.
- 11 McManus E, Sach TH, Levell NJ. An introduction to the methods of decision-analytic modelling used in economic evaluations for Dermatologists. *J Eur Acad Dermatol Venereol* 2019; 33: 1829-1836.
- 12 Petrou S, Gray A. Economic evaluation using decision analytical modelling: design, conduct, analysis, and reporting. BMJ 2011; 342: d1766.
- **13** Sun X, Faunce T. Decision-analytical modelling in health-care economic evaluations. *Eur J Health Econ* 2008; 9: 313-323.
- 14 Wang S, Gum D, Merlin T. Comparing the ICERs in Medicine Reimbursement Submissions to NICE and PBAC - does the

- presence of an explicit threshold affect the ICER proposed? *Value Health* 2018; 21: 938-943.
- **15** Naimark D, Krahn MD, Naglie G, et al. Primer on medical decision analysis: Part 5--Working with Markov processes. *Med Decis Making* 1997; 17: 152-159.
- **16** Caro JJ, Möller J. Advantages and disadvantages of discreteevent simulation for health economic analyses. *Expert Rev Pharmacoecon Outcomes Res* **2016**; **16**: 327-329.
- 17 Caro JJ, Möller J, Karnon J, et al. Discrete event simulation for health technology assessment. United Kingdom: Chapman & Hall, 2015.
- 18 Bojke L, Claxton K, Sculpher M, Palmer S. Characterizing structural uncertainty in decision analytic models: a review and application of methods. *Value Health* 2009; 12: 739-749.
- 19 Briggs AH, Weinstein MC, Fenwick EAL, et al. Model parameter estimation and uncertainty: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force-6. Value Health 2012: 15: 835-842.
- 20 Xiao Y, Hellard ME, Thompson AJ, et al. The cost-effectiveness of universal hepatitis B screening for reaching WHO diagnosis targets in Australia by 2030. Med J Aust 2023; 218: 168-173. https://www.mja.com.au/journal/2023/218/4/cost-effectiveness-universal-hepatitis-b-screening-reaching-who-diagnosis
- 21 Venkataraman P, Neil AL, Mitchell GK, et al. The costeffectiveness of coronary calcium score-guided statin therapy initiation for Australians with family histories of premature coronary artery disease. Med J Aust 2023; 218: 216-222. https:// www.mja.com.au/journal/2023/218/5/cost-effectiveness-coron ary-calcium-score-guided-statin-therapy-initiation
- 22 Husereau D, Drummond M, Augustovski F, et al. Consolidated Health Economic Evaluation Reporting Standards 2022 (CHEERS 2022) statement: updated reporting guidance for health economic evaluations. *BMJ* 2022; 376: e067975.
- 23 Peñaloza Ramos MC, Barton P, Jowett S, Sutton AJ. A systematic review of research guidelines in decision-analytic modeling. Value Health 2015; 18: 512-529.
- 24 Philips Z, Bojke L, Sculpher M, et al. Good practice guidelines for decision-analytic modelling in health technology assessment. *Pharmacoeconomics* 2006; 24: 355-371. ■

Supporting Information

Additional Supporting Information is included with the online version of this article.